

# 6.1 Antiderivatives

---

Standards:

MC11

MC11b

---

---

---

---



# Old Derivatives → Differentiation

Find the derivatives of each.

$$\textcircled{1} f(x) = x^4 + 3x^2 + 6x$$

$$f'(x) = 4x^3 + 6x + 6$$

$$\textcircled{2} y = \ln(2x)$$

$$y' = \frac{1}{2x} (2) = \frac{1}{x}$$

$$\textcircled{3} g(x) = e^{7x^2}$$

$$g'(x) = e^{7x^2} \cdot (14x) \\ = 14xe^{7x^2}$$

$$\textcircled{4} h(x) = 2^{3x^4}$$

$$h'(x) = 2^{3x^4} \cdot \ln(2) \cdot 12x^3 \\ = 12x^3 2^{3x^4} \ln 2$$

$$\textcircled{5} \frac{d}{dx} \sin^{-1}(7x^4) = \frac{1}{\sqrt{1-(7x^4)^2}} \cdot 28x^3 = \frac{28x^3}{\sqrt{1-(7x^4)^2}}$$

$$\textcircled{6} \frac{d}{dx} (x^2y + 2y = 16) = x^2y' + 2y' = 16 \\ [x^2 \cdot y' + y \cdot 2x] + 2y' = 0 \\ x^2y' + 2xy + 2y' = 0 \\ x^2y' + 2y' = -2xy \\ y'(x^2 + 2) = -2xy \\ y' = \frac{-2xy}{x^2 + 2}$$

# New Antiderivatives

## Before



We have been computing derivatives (differentiation) where we start with the position function  $f(x)$  & we "do something" to  $f(x)$  to come up with the derivatives  $f'(x)$ .

## Now

We are going into the "other direction" where we start with  $f(x)$  & we look at another function  $F(x)$  such that  $f(x)$  is the derivative of  $F(x)$  [i.e.  $F'(x) = f(x)$ ].

Let's consider that  $f'(x) = g'(x)$  for all  $x$ , then  $f(x) = g(x) + c$ .

## Example



Conclusion  $g'(x) = 2x$  and  $f'(x) = 2x$ .  $f$  and  $g$  have the same derivative but differ by a constant.

## Definition

If  $F(x)$  is a function such that  $F'(x) = f(x)$ , then  $F(x)$  is called the antiderivative of  $f(x)$ .

Also, if  $F(x)$  is an antiderivative of  $f(x)$ , then  $F(x) + c$  (for some constant  $c$ ) is the most general antiderivative of  $f(x)$ .

[Example]  $x^2 + c$  is the most general antiderivative for  $2x$ .

Question: How can we find antiderivatives of power functions?

We can use what we know about some derivatives.

So here we go:

$$\text{Antiderivatives of } x \Rightarrow \frac{x^2}{2} + C$$

$$x^2 \Rightarrow \frac{x^3}{3} + C$$

$$x^3 \Rightarrow \frac{x^4}{4} + C$$

⋮

$$x^n \Rightarrow \frac{x^{n+1}}{n+1} + C$$

→ power rule for antiderivation

[Examples] Find the most general antiderivatives.

$$\textcircled{1} f(x) = x^5 + 3x^3 + \frac{2}{x^2} - 5e^{-x}$$

Rewrite =  $x^5 + 3x^3 + 2x^{-2} - 5e^{-x}$

$$F(x) = \frac{x^6}{6} + 3\left(\frac{x^4}{4}\right) + 2\left(\frac{x^{-1}}{-1}\right) - 5e^{-x} + C$$

$$= \frac{1}{6}x^6 + \frac{3}{4}x^4 - \frac{2}{x} + 5e^{-x} + C$$

$$\textcircled{2} g(x) = x^4 + 2$$

$$G(x) = \frac{x^5}{5} + 2x + C$$

$$\textcircled{3} h(x) = x^{-2} + \pi$$

$$H(x) = \frac{x^{-1}}{-1} + \pi x + C$$

$$= -\frac{1}{x} + \pi x + C$$

$$\textcircled{4} j(x) = 2x - \frac{3}{x^4}$$

Rewrite =  $2x - 3x^{-4}$

$$J(x) = 2\left(\frac{x^2}{2}\right) - 3\left(\frac{x^{-3}}{-3}\right) + C$$

$$= x^2 + \frac{1}{x^3} + C$$

WARNING: We do not have analogies of product, quotient or chain rules for antidifferentiation... (yet!)

$$\textcircled{5} f(x) = \frac{x^3 + x^2 + x}{x}$$

Rewrite =  $\frac{x^3}{x} + \frac{x^2}{x} + \frac{x}{x} = x^2 + x + 1$

$$F(x) = \frac{x^3}{3} + \frac{x^2}{2} + x + C$$

Remember:

$$\frac{d^2}{dt^2} s(t) = \frac{d}{dt} v(t) = a(t)$$

↑  
position  
function

↑  
velocity  
function

↑  
acceleration  
function

$f(x)$  = position

$f'(x)$  = velocity

$f''(x)$  = acceleration

Integration → the process of antidifferentiation of a function.

## Definition

Given a function,  $F(x)$ , an antiderivative of  $f(x)$  is any function  $F(x)$  such that  $F'(x) = f(x)$ .

If  $F(x)$  is any antiderivative of  $f(x)$ , then the most general antiderivative of  $f(x)$  is called an indefinite integral and denoted as:

$$\int f(x) dx = F(x) + C$$

Integral symbol      integrand      "x" is the integration variable      antiderivative

[Examples] Integrate the functions.

$$\textcircled{1} \int x^4 + 3x - 9 dx = \frac{x^5}{5} + \frac{3x^2}{2} - 9x + C$$

$$\textcircled{2} \int x^2 - 2x - 5 dx = \frac{x^3}{3} - \frac{2x^2}{2} - 5x + C$$
$$= \frac{1}{3}x^3 - x^2 - 5x + C$$

$$\textcircled{3} \int 2 - 5x^2 dx = 2x - \frac{5x^3}{3} + C$$

Why is  $\int f(x) dx = F(x) + C$  called an indefinite integral?

The reason is because we are just finding the antiderivative, not evaluating the antiderivative.

What about other types of functions?

Antiderivatives for Trig Functions:

$$\begin{aligned} \int \sin x dx &= -\cos x + C & \int \csc x \cot x dx &= -\csc x + C \\ \int \cos x dx &= \sin x + C & \int \sec x \tan x dx &= \sec x + C \\ \int \sec^2 x dx &= \tan x + C & \int \csc^2 x dx &= -\cot x + C \\ \int \tan x dx &= \ln|\sec x| + C & \int \cot x dx &= \ln|\sin x| + C \\ \int \sec x dx &= \ln|\sec x + \tan x| + C & \int \csc x dx &= \ln|\csc x - \cot x| + C \end{aligned}$$

Antiderivatives for Inverse Trig Functions:

$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + C \quad \int \frac{1}{1+x^2} dx = \tan^{-1} x + C$$

Antiderivatives for Polynomials:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C$$



## Antiderivatives for Exponential Rules/Logarithmic Rules

$$\int e^x dx = e^x + c$$

$$\int \frac{1}{x} dx = \ln|x| + c$$

## Position Function Problems

Let's say  $f'(x) = x^3$  and  $f(0) = 4$ . What is the position function.

$$f'(x) = x^3$$

$$f(x) = \frac{x^4}{4} + C$$

Now we know that  $f(0) = 4$ , so...  $f(0) = \frac{(0)^4}{4} + C = 4$   
 $C = 4$

The position function is  $f(x) = \frac{x^4}{4} + 4$  when  $f(0) = 4$ .

[Example] Find the position function.

$$\textcircled{1} f'(t) = 2t + 9t^2, \quad f(1) = 2$$

$$f(t) = \frac{2t^2}{2} + \frac{9t^3}{3} + C$$
$$= t^2 + 3t^3 + C$$

$$\text{Since } f(1) = 2, \quad f(1) = (1)^2 + 3(1)^2 + C = 2$$
$$= 1 + 3 + C = 2$$
$$= 4 + C = 2$$
$$C = -2$$

Therefore when  $f(1) = 2$ ,  $f(t) = t^2 + 3t^3 - 2$ .

$$\textcircled{2} f''(x) = x, \quad f(0) = -3, \quad f'(0) = 2$$

$$f''(x) = x$$

$$f'(0) = \frac{0^2}{2} + c = 2$$

$$f'(x) = \frac{x^2}{2} + c$$

$$c = 2$$

$$f(x) = \frac{x^3}{6} + cx + d =$$

$$f(0) = \frac{0^3}{6} + 2(0) + d = -3$$

$$d = -3$$

Therefore,  $f(x) = \frac{x^3}{6} + 2x - 3$  when  $f(0) = -3$  and  $f'(0) = 2$ .